Grief and Loss Resources

WEB SITES:

Al-Anon Family Groups District 5: http://www.afgdistrict5.org/

Compassion Books: http://www.compassionbooks.com/store/

American Society of Suicidology: http://www.suicidology.org/

Center for Loss and Life Transition: s http://www.centerforloss.com/

Compassionate Friends: Supporting a Family after a Child Dies:

http://www.compassionatefriends.org/

Crisis, Grief & Healing: http://www.webhealing.com/

Griefnet: http://www.griefnet.org/

Hospice Net: http://www.hospicenet.org/

"Denial to Acceptance: The Stages of Grief in Addiction and Recovery"

http://www.addictiontreatmentmagazine.com/recovery/recovery-tips/denial-to-acceptance-the-stages-of-grief-in-addiction-and-recovery/

BOOKS:

Rudman, M.K., Gagne K. D., Bernstein, J.E., <u>Books to Help a Child Cope with Separation and Loss:</u> An Annotated Bibliography 4th ed., 1994.

Fitzgerald, Helen, The Grieving Child, 1992.

Jarratt, C.J., Helping Children Cope with Separation and Loss, Revised Edition, 1994.

Worden, J. William, Grief Counseling & Grief Therapy, 2008.

Viorst, Judith, Necessary Losses, 2002.

Lynn, J. and Harrold, J. (1999). <u>Handbook for Mortals: Guidance for People Facing Serious Illness</u>, p.41.

Abbot, A., Alcohol, Tobacco, and Other Drugs. NASW, 2010.

Self-Care

Adapt your priorities and expectations

- Each day plan what is most important to you
- Alter schedule to plan for difficult tasks
- Make yourself a priority
- Accept less than perfection from self and others
- Know that others are also working with this client

Adjust your attitude

- Talk with others who are doing similar work
- Open up your outlook often we see what we look for
- Stop "awefulizing"
- Acknowledge value of "small acts" and witnessing, listening, validating, caring

Act out of caring and respect for you

- Balance "being" and "doing"
- Be vulnerable and human
- Reward and reinforce yourself and those around you
- Say "no" sometimes
- Take your turn

Acclaim your achievements and strengths

- Choose a "strength for the day" to focus on and enjoy
- Keep a journal of your accomplishments and experiences
- Recognize your skill and give yourself credit
- Acknowledge and save compliments

Acknowledge your own needs

- Accept that you have needs physical, emotional, spiritual
- Recognize limitations as acceptable
- Try to be aware of your "buttons" and loss history
- Talk about your feelings of helplessness, vulnerability, etc.
- Take breaks, meditate, relaxation, imagery throughout the day

Activate your support system

- Know where your supports are
- Develop a nurture network
- Find a way to recognize, replenish, reward yourself each day
- Know what your personal warning signs are and confront first signs of stress
- Use journaling, art work, hobbies, exercise, vacations, cry
- Network with other team members
- Go to conferences and professional meetings

Courtesy Debbie Mattison, LMSW, UM School of Social Work

Janice Firn, LMSW Clinical Social Worker A comparison of characteristics of both helpers and rescuers:

The Helper

- 1. Listens for request
- 2. Presents offer
- 3. Gives only what is needed
- 4. Checks periodically with person
- 5. Checks results
 - a. Functioning better?
 - b. Meeting goals?
 - c. Solving problems independently?
 - d. Using suggestions successfully?

The Rescuer

- 1. Gives when not asked
- 2. Neglects to find out if offer is welcome
- 3. Gives help more and longer than needed
- 4. Omits feedback
- 5. Doesn't check results and feels good when accepted, bad when turned down
- 6. Does the greater share of the talking

Excerpt from Wellness Workbook by Regina S. Ryan & John W. Travis, MD.

Being responsible TO others is FREEING to self and others

Being responsible FOR others is CONTROLING for self and others

- 1. Listening, empathy
- 2. Separate from the other at least a minimal degree of objectivity
- 3. Personal (focus on the other as a person rather than a "thing" or object
- 4. Loves
- Congruent/Genuine, "hears" body, deals with feelings constructively (especially anger), confronts constructively, constructive feedback (positive & negative)
- 6. Respects the other person
 - a. Provides freedom to choose (considers options)
 - b. Allows freedom to succeed or "fail"
 - c. Confident of other

- 1. Telling, sympathy, pity; reassurance only, positive feedback only
- 2. Immersed or over-involved emotionally or coldly detached
- 3. Depersonalizing, stereotyping, labeling (the "case", the problem in room 303, the "sociopath", the "discipline problem")
- 4. Smothers
- 5. Hostility (active or passive), Incongruent/Phony, Denies or ignores body, denies or distorts feelings, avoids confrontation
- 6. Lacks confidence
 - a. Only my answer or decision is OK (persuade, "sweetly" coerce, bribe, blame, increase guilt, withdraws affection, etc.)
 - b. Stays awake nights, worries, takes problem home
 - c. Failure is tragic
- 7. Super-helper, savior, must save everyone, perfectionist, super-giver

7. Helper can fail

Excerpt from Pancrazio & Zeller 1982

Rescuers Checklist

Completing the checklist can help you become aware of the ways you may be rescuing people without realizing it. It is taken, with permission, from the Transactional Checklist. Mark each of the statements below as it applies to you: 0 = seldom or never; 1 = sometimes or occasionally; and 2 = frequently. X = significant others in your life such as a spouse, boss, parents, friend, or colleague.

 _ 1. Is it hard for you to take time for yourself and have fun?
 2. Do you supply words for X when he/she hesitates?
 _ 3. Do you set limits for yourself that you then exceed?
 4. Do you believe that you are responsible for making (keeping) X happy?
 _ 5. Do you enjoy lending a shoulder for X to "cry" on?
 6. Do you believe that X is not sufficiently grateful for your help?
 7. Do you take care of X more than you take care of yourself?
 8. Do you find yourself interrupting when X is talking?
 9. Do you watch for clues for ways to be helpful to X?
 _ 10. Do you make excuses, openly or mentally, for X?
 _ 11. Do you do more than your share, that is, work harder than X?
 _ 12. When X is unsure or uncomfortable about doing something do you do it for X?
 _ 13. Do you give up doing things because X wouldn't like it?
 _ 14. Do you find yourself thinking that you really know what is best for X?
 _ 15. Do you think X would have grave difficulty getting along without you?
 _ 16. Do you use the word "we" and then find out you don't have X's consent?
 _ 17. Do you stop yourself by thinking X will feel badly if you for or do something?
 _ 18. Is it hard for you not to respond to anyone who seems hurting or needing help?
 _ 19. Do you find yourself being resented when you are only trying to help?
 20. Do you find yourself giving advice that is not welcome or accepted?
 _ Total: More than 10 points – rescuing is possible, more than 20 points – rescuing is probable.

Children & Grief

Children's experience of loss and grief can differ from an adult's experience. Knowing how a child's developmental stage affects his/her understanding of and ability to cope with loss is important. This knowledge will help you provide the most meaningful and effective support to your child. Speaking with your child's health care provider regarding his/her response to loss can be a helpful place to start. Pediatricians can assess whether a child's response is normal or if professional support/counseling is needed. Additionally, it can be beneficial to involve the school counselor or social worker; they can also assess for coping and identify support/counseling resources. Below is a list of resources for helping adults to better understand children's grief, and options for how to support children through the grieving process.

American Academy of Child and Adolescent Psychiatry

http://www.aacap.org/cs/forFamilies

Arbor Hospice

http://www.arborhospice.org/we-can-help/grief-support-services

Barr-Harris Children's Grief Center

http://www.barrharris.org/

The Center for Grieving Children, Teens, and Families

http://grievingchildren.org/

The Children's Room

http://childrensroom.org/

Ele's Place

http://www.elesplace.org/

The Dougy Center

http://www.dougy.org/grief-resources/how-to-help-a-grieving-child/

Hospice Net

http://www.hospicenet.org/html/child.html

Michigan Mental Health Networker, Washtenaw County Child & Adolescent Services

http://www.mhweb.org/index.html

http://www.mhweb.org/washtenaw/child_index.html

Avoiding the Clichés of Grief

Try open ended statements. Try to avoid telling people what they "should" or "should not" be feeling, doing, etc. Do not assume to know what someone is going through or experiencing.

1. Cliché: "You must be strong for your children (spouse, relatives, friends, etc)." Instead, try: Why not share your feelings with your children? Perhaps you can lean on one another and help support each other.

2. Cliché: "You've got to get hold of yourself."

Instead, try: "It must be so hard to keep going when you're hurting so much."

3. Cliché: "You are holding up so well."

Instead, try: "Would it help to talk about how you're feeling?"

4. Cliché: "Time will heal."

Instead, try: "You must feel as if this pain will never end."

5. Cliché: "You're young, and you will be able to make a new life for yourself."

Instead, try: "You must miss your loved one and the life you had together; I do, too."

What to Say

What *Not* to Say

I'm sorry. I understand how you feel.

I'm sad for you. Death was a blessing.

How are you doing with all this? It was God's will.

I don't know why it happened. It all happened for the best.

What can I do for you? You're still young.

I'm here and I want to listen. You have your whole life ahead of you.

Please tell me what you are feeling. You can have other children.

This must be hard for you. You can always remarry.

What's the hardest part for you? Call me when I can help.

I'll call tomorrow. Something good will come of this. You must really be hurting. At least you have another child.

It isn't fair, is it? She/he led a full life.

You must really feel angry. It's time to put it behind you.

Take all the time you need. Be strong!

Courtesy, Archdiocese of Omaha, Family Life Office

24. Change in social activities	29
25. Change in eating habits	28
26. Chronic car trouble	26
27. Change in number of family get-togethers	26
28. Too many missed classes	25
29. Changing colleges	24
30. Dropping more than one class	23
31. Minor traffic violations	20

Total	Stress	Score	

Score Interpretation:

Researchers determined that if your total score is:

300 or more - statistically you stand an almost 80 percent chance of getting sick in the near future.

150 to 299 - you have a 50-50 chance of experiencing a serious health change within two years.

149 or less - you have about a 30 percent chance of a serious health change.

This scale indicates that change in one's life requires an effort to adapt and then an effort to regain stability. Stress is a natural byproduct of adapting and then regaining internal homeostasis. Take note that this assessment considers only the events that occur, not individual perception of these events in life. Perception is a critical part of the ultimate stress experience, so while the Student Stress Scale has value in increasing awareness of potential stress-producing events, ultimately individual perception of the event is an important variable.

The Student Stress Scale

The Student Stress Scale focuses on *events* that may occur in the life of a student to offer you a different perspective for evaluating stress. The Student Stress Scale is an adaptation for college students of the Life Events Scale developed originally by Holmes and Rahe. This popular stress assessment measured the amount of change, using Life Change Units, a person was required to adapt to in the previous year. It was designed to predict the likelihood of disease and illness following exposure to stressful life events. Each life event is given a score that indicates the amount of readjustment a person has to make as a result of change. Some studies have found that people with serious illnesses tend to have higher scores on similar assessments.

For each event that occurred in your life within the past year, record the corresponding score. If an event occurred more than once, multiply the score for that event by the number of times the event occurred and record that score. Total all the scores.

Life Event	Mean Value
1. Death of a close family member	100
2. Death of a close friend	73
3. Divorce of parents	65
4. Jail term	63
5. Major personal injury or illness	63
6. Marriage	58
7. Getting fired from a job	50
8. Failing an important course	47
9. Change in the health of a family member	45
10. Pregnancy	45
11. Sex problems	44
12. Serious argument with a close friend	40
13. Change in financial status	39
14. Change of academic major	39
15. Trouble with parents	39
16. New girlfriend or boyfriend	37
17. Increase in workload at school	37
18. Outstanding personal achievement	36
19. First quarter/semester in college	36
20. Change in living conditions	31
21. Serious argument with an instructor	30
22. Getting lower grades than expected	29
23. Change in sleeping habits	29

Life Events Scale

This stress assessment measures the amount of change, using Life Change Units, a person experienced and adjusted to in the previous 12 months. It was designed to predict the likelihood of disease and illness following exposure to stressful life events. Each life event is given a score that indicates the amount of readjustment a person has to make as a result of the event. Not all of the events in the scale are necessarily negative events.

This scale indicates that change in one's life requires an effort to adapt and then an effort to regain stability. Stress and feelings of loss are natural by-products of adapting and trying to regain homeostasis. This assessment considers only the events that occurred, not individual perception of these events in life. Perception is a key part of the total grief and loss experience, so while the Life Events Scale has value in increasing awareness of potential losses and life changing or stressful events, an individual's perception of the event is an important variable which needs to be considered in the overall assessment. For a more complete picture of how loss has affected a person's life consider all the dimensions of health: physical, mental, emotional, spiritual, and social.

Directions

For each event that occurred in your life within the past year, record the corresponding score. If an event occurred more than once, multiply the score for that event by the number of times the event occurred and record that score. Total all the scores:

- Score of 300+: At risk of illness.
- Score of 150-299+: Risk of illness is moderate (reduced by 30% from the above risk).
- Score 150-: Only have a slight risk of illness.

A modified scale has also been developed for students (teenagers and university aged young adults). This scale is included in the handout. Similar to the adult scale, life events are totaled and provide a rough estimate of how life changes may affect health.

Adapted from Holmes-Rahe Social Readjustment Rating Scale. Journal of Psychosomatic Research, (1967). Vol. 11, pp. 213-218.

Grief: Suggestions for Finding a Support Group or Therapist

Receiving support from others who have experienced a loss along with receiving professional support can help us cope with a loss. It may seem a daunting task to find the appropriate support group or therapist. Many local religious leaders or places of worship, hospitals, treatment centers, hospices, and funeral homes have lists of support groups. Even if you or your loved one was not a patient or client of these organizations most are open to community members and are willing to assist you with identifying support options.

Most types of loss have specific web sites (ex: Suicide, Miscarriage, Substance Use); these sites often have helpful suggestions for support groups, agencies, and networks. Ask friends, family or peers who are also experiencing the same loss for recommendations of groups or therapists.

Another way to identify therapists in your area is to contact your health care provider (doctor/clinic) or health insurance provider (ex: Blue Cross Blue Shield, BCN, HAP, Medicaid HMO, etc.). Under the Affordable Care Act insurance companies are required to cover mental health services. However, the number and type of visits covered will vary by plan. Contact your plan for more information.

In the Ann Arbor area there are several community agencies and groups that may be able to assist you with finding a group or therapist, below are a few of these agencies:

Washtenaw County Mental Health: 734-544-3050 or 1-800-440-7548 (24 hours) St. Joseph's Mercy Behavioral Health: 734-786-2301 or 1-800-289-0014 University of Michigan Health System Psychiatry: 734-764-0231 or 1-800-525-5188

Ann Arbor Area Therapeutic Resources:

http://www.therapeuticresources.com/supportmichigan.html

Arbor Hospice: http://www.arborhospice.org/we-can-help/grief-support-services

Community Support and Treatment Services:

http://www.ewashtenaw.org/government/departments/community_mental_health/programs-and-services

Ele's Place: http://www.elesplace.org/ - Children Groups & Support

Grief Net:

http://griefnet.org/support/groups.html - Adult Groups http://kidsaid.com/ - Children's Groups

Michigan Mental Health Networker, Washtenaw County

http://www.mhweb.org/washtenaw/selfhelp_index.html http://www.mhweb.org/washtenaw/therapist_indx.html

ACA information: https://www.healthcare.gov/do-marketplace-insurance-plans-cover-mental-health-and-substance-abuse-services/

Janice Firn, LMSW Clinical Social Worker